


Integrating outcrop and subsurface data to assess the temporal evolution of a submarine channel–levee system

Emma A. Morris, David M. Hodgson, Stephen Flint, Rufus L. Brunt, Stefan M. Luthi, and Yolanda Kolenberg

AAPG Bulletin, v. 100, no. 11 (November 2016), pp. 1663–1691



Copyright ©2016. The American Association of Petroleum Geologists. All rights reserved. Green Open Access. This paper is published under the terms of the CC-BY license.

Table A. Channel Axis





FMS = formation microscanner.

Table B. Channel Margin

(B)	<i>Outcrop photographs Facies associations</i>	<i>Core photographs Facies associations</i>	
Channel Margin			
	<i>FMS image log facies associations</i>		
			




FMS = formation microscanner.

Table C. Proximal Internal Levee

(C)	<i>Outcrop photographs Facies associations</i>	<i>Core photographs Facies associations</i>	
Proximal Internal Levee			
	<i>FMS image log facies associations</i>		
			




FMS = formation microscanner.

Table D. Distal Internal Levee

(D)	<i>Outcrop photographs Facies associations</i>	<i>Core photographs Facies associations</i>
Distal Internal Levee		
	<i>FMS image log facies associations</i>	
		

FMS = formation microscanner.

Table E. Proximal External Levee

(E) Proximal External Levee	<i>Outcrop photographs Facies associations</i>	<i>Core photographs Facies associations</i>	
			
	<i>FMS image log facies associations</i>		
			

FMS = formation microscanner.

